

Attachment B

how to read a drug label

Who

specific about your pain.

View all profiles

Print this page

Email this page

Most over-the-counter medicines have a Drug Facts label to help you select the right product and use it correctly.

Roll over the numbers on the sample label below for a detailed explanation.

1	Drug Facts Active ingredient (in each caplet) Ibuprofen 200 mg (NSAID) *nonsteroidal anti-inflammatory drug	Purpose Pain reliever/ fever reducer				
2	Uses Temporarily relieves minor aches and pains due to: • headache • muscle aches • minor pain of arthritis • toothache • backache • the common cold • menstrual cramps • temporarily reduces fever					
3	Warnings Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include: • hives • facial swelling • asthma (wheezing) • shock • skin reddening • rash • blisters If an allergic reaction occurs, stop use and seek medical help right away. Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is higher if you: • are age 60 or older • have had stomach ulcers or bleeding problems • take a blood thinning (anticoagulant) or steroid drug • take other drugs containing an NSAID (aspirin, ibuprofen, naproxen, or others) • have 3 or more alcoholic drinks every day while • take more or for a longer time than directed Do not use • if you have ever had an allergic reaction to any other pain reliever/fever reducer • right before or after heart surgery Ask a doctor before use if you have: • problems or serious side effects from taking pain relievers or fever reducers • stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain • ulcers • bleeding problems • high blood pressure • heart or kidney disease • taken a diuretic • reached age 60 or older Ask a doctor or pharmacist before use if you are: • taking any other drug containing an NSAID (prescription or nonprescription) • taking a blood thinning (anticoagulant) or steroid drug • under a doctor's care for any serious condition • taking any other drug					
	Drug Facts (continued) When using this product • take with food or milk if stomach upset occurs • long term continuous use may increase the risk of heart attack or stroke Stop use and ask a doctor if: • you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding. • pain gets worse or lasts more than 10 days • fever gets worse or lasts more than 3 days • stomach pain or upset gets worse or lasts • redness or swelling is present in the painful area • any new symptoms appear If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)					
	Directions • do not take more than directed • the smallest effective dose should be used • do not take longer than 10 days, unless directed by a doctor (see Warnings) <table border="1"> <tr> <td data-bbox="714 1155 812 1344">adults and children 12 years and older</td> <td data-bbox="812 1155 1102 1344"> • take 1 caplet every 4 to 6 hours while symptoms persist • if pain or fever does not respond to 1 caplet, 2 caplets may be used • do not exceed 6 caplets in 24 hours, unless directed by a doctor • ask a doctor </td> </tr> <tr> <td data-bbox="714 1344 812 1491">children under 12 years</td> <td data-bbox="812 1344 1102 1491"></td> </tr> </table>		adults and children 12 years and older	• take 1 caplet every 4 to 6 hours while symptoms persist • if pain or fever does not respond to 1 caplet, 2 caplets may be used • do not exceed 6 caplets in 24 hours, unless directed by a doctor • ask a doctor	children under 12 years	
adults and children 12 years and older	• take 1 caplet every 4 to 6 hours while symptoms persist • if pain or fever does not respond to 1 caplet, 2 caplets may be used • do not exceed 6 caplets in 24 hours, unless directed by a doctor • ask a doctor					
children under 12 years						
	Other information • store between 20 - 25°C (68 - 77°F) • do not use if neck wrap or foil inner seal imprinted "Safety Sealer" is broken or missing • see end panel for lot number and expiration date					
	Inactive ingredients carnauba wax, corn starch, FD&C yellow #6, hypromellose, iron oxide, polydextrose, polyethylene glycol, silicon dioxide, stearic acid, titanium dioxide Questions or comments? Call 1-877-893-3463 (English) or 1-888-486-8746 (Spanish) weekdays 9:00 AM to 4:30 PM EST					



Wyeth Consumer Healthcare | Cor

Advil The Every Pain

About New Advil® Labeling

- ☐ Advil Home
- ☐ Adult Products
 - ☐ Advil
 - ☐ Advil PM
 - ☐ Advil Migraine
 - ☐ Advil Cold & Sinus
 - ☐ Advil Allergy Sinus
 - ☐ Advil Multi-Symptom Cold
- ☐ Children's Products
- ☐ Pain Free Living
- ☐ FAQ
- ☐ Buy Online
- ☐ Find A Store Near You
- ☐ Wyeth Special Offers

This is the most current labeling information, and may differ from labels on product packaging. If there are any differences between this website labeling and product package labeling, this website labeling should be regarded as the most current.

The FDA has approved enhanced warning labels for all over the counter non-steroidal anti-inflammatory drug analgesics (NSAID), including all Advil® products. These new labels will provide consumers with more information about potential cardiovascular, gastrointestinal and allergic risks.

Products with revised labels should be in stores by late 2006. In the meantime, we have posted the revised label on this Web site. We continue to work with the FDA to ensure the effective labeling of our products.

Wyeth Consumer Healthcare stands firmly behind the safety and effectiveness of Advil®. However, we encourage all consumers to read the new, enhanced Advil® labeling, with special attention to the Warnings. All drugs, including those without a prescription, must be used according to their directions to be safe and effective.

Advil® Liqui-Gels®

Drug Facts

Active Ingredient (in each capsule)

Solubilized ibuprofen equal to 200 mg ibuprofen (NSAID)

(present as the free acid and potassium salt)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/Fever reducer

Uses

temporarily relieves minor aches and pains due to:

headache toothache backache menstrual cramps
the common cold muscular aches minor pain of arthritis

temporarily reduces fever

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

hives facial swelling asthma (wheezing)
shock skin reddening rash blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains a nonsteroidal anti-inflammatory drug (NSAID), which may cause stomach bleeding. The chance is

higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing an NSAID [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Do not use

If you have ever had an allergic reaction to any other pain reliever/fever reducer right before or after heart surgery

Ask a doctor before use if you have

- problems or serious side effects from taking pain relievers or fever reducers
- stomach problems that last or come back, such as heartburn, upset stomach, or stomach pain
- ulcers
- bleeding problems
- high blood pressure
- heart or kidney disease
- taken a diuretic
- reached age 60 or older

Ask a doctor or pharmacist before use if you are

- taking any other drug containing an NSAID (prescription or nonprescription)
- taking a blood thinning (anticoagulant) or steroid drug
- under a doctor's care for any serious condition
- taking aspirin for heart attack or stroke, because ibuprofen may decrease this benefit of aspirin
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs
- long term continuous use may increase the risk of heart attack or stroke

Stop use and ask a doctor if

- you feel faint, vomit blood, or have bloody or black stools. These are signs of stomach bleeding.
- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- stomach pain or upset gets worse or lasts
- redness or swelling is present in the painful area
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use ibuprofen during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than directed**
- the smallest effective dose should be used**
- do not take longer than 10 days, unless directed by a doctor (see Warnings)
- adults and children 12 years and over: take 1 capsule every 4 to 6 hours while symptoms persist
- if pain or fever does not respond to 1 capsule, 2 capsules may be used
- do not exceed 6 capsules in 24 hours, unless directed by a doctor
- children under 12 years: ask a doctor

Other information

- each capsule contains:** potassium 20 mg
- read all warnings and directions before use. Keep carton.

store at 20-25°C (68-77°F)
avoid excessive heat above 40°C (104°F)

Inactive ingredients

FD&C green no. 3, gelatin, light mineral oil, pharmaceutical ink, polyethylene glycol, potassium hydroxide, purified water, sorbitans, sorbitol

Questions or comments? call toll-free 1-800-88-ADVIL

[► Back to Product Page](#)

© 2008 Wyeth Consumer Healthcare [Privacy](#) · [Terms & Conditions](#) · [Site Map](#)

This site is intended for residents of the [United States](#).

Attachment D

RECEIVED
CENTRAL FAX CENTER

APR 18 2008

[About Us](#)
[Cataloged List](#)
[Alphabetic List](#)
[What's New](#)
[Contact](#)

Upper level

Acesulfame K

Aspartame

Calcium chloride

Erythorbic acid

DL-Lactic acid

L-Lactic acid

Potassium sorbate


Sodium benzoate

Sodium erythorbate

Sorbic acid

Sodium Benzoate

Identification

Synonyms: Isoascorbic Acid Sodium Salt; Sodium Isoascorbate**CAS No.:** 532-32-1**Molecular Weight:** 144.10**Molecular Formula:** C₇H₅NaO₂**Structural Formula:**

Application

sodium benzoate is applied as an antioxidant used in meat, poultry, soft drink.

Specification

Complied with BP98/FCC IV, Kosher Certified

Item	Specification
sodium benzoate purity	99.0%min.
moisture	1.5% max.
chlorides as Cl	200ppm max.
total chlorides	300ppm max.
heavy metals as Pb	0.001% max.
arsenic as As	0.0003% max.
alkalinity	0.2ml/g max.
color of solution	Y6
appearance	white crystalline powder or granule

Standard Packing

25kgs net/woven bags with plastic inner bags, or Kraft bag

Risks and Safety Info

Safe for land, air and water transportation

Availability

Attachment E

Ibuprofen was made available under prescription in the United Kingdom in 1969, and in the United States in 1974. In the years since, the good tolerability profile along with extensive experience in the community (otherwise known as Phase IV trials), has resulted in the rescheduling of small packs of ibuprofen to allow availability over-the-counter in pharmacies worldwide, and indeed in supermarkets and other general retailers. For some time, there has been a limit on the amount that can be bought over the counter in a single transaction in the UK; this being 2 packs of 16 x 200 mg or 1 pack of 8 or 16 x 400 mg, the latter being far less common for over-the-counter sales. In the United States, the Food and Drug Administration approved it for over-the-counter use in 1984. The wider availability has meant that ibuprofen is now almost as commonly used as aspirin and paracetamol (acetaminophen).^[citation needed] In other countries, such as Spain, higher dosages of 600 mg are available. Mexico's doses go up to 800mg per pill.